

UCRL- 101082  
PREPRINT

OXYGEN VACANCY ORDERING IN  $\text{YBa}_2\text{Cu}_3\text{O}_{7-y}$

P.A. STERNE  
L.T. WILLE

THIS PAPER WAS PREPARED FOR SUBMITTAL TO  
INTERNATIONAL M<sup>2</sup>S-HTSC CONFERENCE  
Stanford, CA  
July 23-28, 1989

June, 1989

Lawrence  
Livermore  
National  
Laboratory

This is a preprint of a paper intended for publication in a journal or proceedings. Since changes may be made before publication, this preprint is made available with the understanding that it will not be cited or reproduced without the permission of the author.

COMMUNICATION COPY  
SUBJECT TO RECALL  
TWO WEEKS

## DISCLAIMER

This document was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor the University of California nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial products, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement recommendation, or favoring of the United States Government or the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or the University of California, and shall not be used for advertising or product endorsement purposes.

OXYGEN VACANCY ORDERING IN  $\text{YBa}_2\text{Cu}_3\text{O}_{7-y}$

P. A. STERNE

Lawrence Livermore National Laboratory, Livermore, CA 94550

L. T. WILLE

Physics Department, Florida Atlantic University, Boca Raton, FL 33431

First principles total energy calculations have been performed for  $\text{YBa}_2\text{Cu}_3\text{O}_x$  with  $x$  varying from 6.0 to 7.5. The results of these calculations have been used to determine the effective pair interactions for the oxygen ordering in the basal plane. The phase diagram calculated with these "first principles" pair interactions is in very good agreement with experiment. Comparison of the total energies for the  $\text{O}_8$ ,  $\text{O}_{8.5}$  and  $\text{O}_7$  structures shows that the double-cell phase is thermodynamically stable.

The problem of oxygen ordering in the basal plane of  $\text{YBa}_2\text{Cu}_3\text{O}_{7-y}$  is an important one which has received a great deal of attention recently. Experimentally<sup>1</sup>, it is clear that the tetragonal to orthorhombic structural transition is associated with an ordering of the oxygen atoms into Cu-O chains. Theoretically, de Fontaine, Wille and co-workers<sup>2</sup> have proposed a simple Ising like model to describe the interactions between the oxygen atoms and have shown that, for a particular range of parameters, their model gives rise to a phase diagram in qualitative agreement with experiment. In contrast to this method which is based on localized effective pair interactions, Khachaturyan and Morris<sup>3</sup> have suggested an alternative approach where long range interactions produce the observed ordering. They conclude that the double-cell phase of  $\text{YBa}_2\text{Cu}_3\text{O}_{8.5}$ , with parallel Cu-O chains running through every second copper atom in the basal plane, is thermodynamically unstable, in disagreement with the conclusions of de Fontaine et al.

One way to resolve this controversy is to calculate the total energies of the structures using a first principles electronic structure method and see if the double-cell phase is

higher or lower in energy than the average of the  $\text{O}_8$  and  $\text{O}_7$  phases. We have performed these calculations using the LMT0 method<sup>4</sup>. The double cell phase was found to be more stable by 4.6mRy/O atom in the basal plane, and will clearly not decompose into a mixed  $\text{O}_8/\text{O}_7$  phase.

The model proposed by de Fontaine et al<sup>2,5</sup> is an Ising like model with nearest and next nearest neighbor interactions. There are two different next nearest sites in the basal plane, one joined by a Cu atom, the other with no intervening atoms, so it is assumed that these two interactions are different, giving rise to an asymmetric next nearest neighbor Ising model described by three interactions:  $V_1$ , the nearest neighbor interaction,  $V_2$ , the second-neighbor interaction mediated by the Cu atom and  $V_3$ , the second-neighbor interaction with no intervening atoms. Wille et al<sup>2</sup> have shown that for an appropriate choice of parameters, this model can produce a phase diagram in qualitative agreement with experiment. The choice of parameters is very important, since the phase diagram can be qualitatively altered by choosing different values.

We have calculated these parameters from first principles by computing the total energies

for a number of different structures and using the Connolly Williams<sup>8</sup> method to extract the interaction energies. Care must be taken in deriving these parameters from the electronic structure calculations, since the total energies are of order thousands of Rydbergs, while the interaction parameters are only a few mRy. The following values were obtained:

$$\begin{aligned} V_1 &= 8.9 \text{ mRy} & V_2 &= -2.4 \text{ mRy} \\ V_3 &= 1.1 \text{ mRy} \end{aligned}$$

The phase diagram in figure 1 has been calculated from these parameters using the cluster variation method<sup>7</sup>. Agreement with experiment for the orthorhombic to tetragonal transition is excellent, considering that this is a first principles calculation with no fitting to experiment. The double-cell phase, labelled Ortho II, is clearly seen around  $O_{6.5}$ . The stability of this phase can be readily




FIGURE 1

Phase diagram for  $YBa_2Cu_3O_{7-y}$  calculated from the pair interactions derived from the total energy calculations. The diamonds<sup>8</sup> and squares<sup>9</sup> are experimental data points.

understood in terms of the calculated parameters: the negative  $V_2$  term indicates that it is energetically favorable to form Cu-O chains (or, equivalently, Cu-vacancy ( $\square$ ) chains), as seen in both  $YBa_2Cu_3O_7$  and  $YBa_2Cu_3O_{6.8}$ . The repulsive  $V_3$  term suggests that the system would favor a state where the O-O and  $\square$ - $\square$  repulsive interactions are minimized, which is the case for the double cell phase.

#### ACKNOWLEDGEMENTS

This work was performed under the auspices of the U.S. Department of Energy by the Lawrence Livermore National Laboratory under contract W-7405-ENG-48. Part of the work was done while one of the authors (PAS) was at the University of Maryland where it was supported by ONR contract N00014-86-K-0266. LTW acknowledges financial support from DARPA grant number MDA-972-88-J-1006. Calculations were carried out at the Pittsburgh Supercomputer Center and the NMFE Computer Center.

#### REFERENCES

1. R. Beyers and T.M. Shaw, Solid State Physics 42 (1989) 135
2. D de Fontaine, L.T. Wille and S.C. Moss, Phys. Rev. B 36 (1987) 5709; L.T. Wille and D. de Fontaine, Phys. Rev. B 37 (1988) 2227; L.T. Wille, A. Berera and D. de Fontaine, Phys. Rev. Lett. 60 (1988) 1065
3. A.G. Khachaturyan and J.W. Morris Jr., Phys. Rev. Lett. 59 (1987) 2776; Phys. Rev. Lett. 61 (1988) 215
4. P.A. Sterne and L.T. Wille, to be published
5. L.T. Wille, Phase Transitions, in print
6. J.W.D. Connolly and A.R. Williams, Phys. Rev. B 27 (1983) 5169
7. R. Kikuchi, Phys. Rev. 81 (1951) 988
8. Y. Kubo and H Igarashi, Phys. Rev. B 39 (1989) 725
9. E.D. Specht et al, Phys. Rev. B 37 (1988) 7246